

Matter-gravity scattering in the presence of spontaneous Lorentz violation

R. V. Maluf,^{1,*} Victor Santos,^{1,†} W. T. Cruz,^{2,‡} and C. A. S. Almeida^{1,§}

¹*Universidade Federal do Ceará (UFC),
Departamento de Física, Campus do Pici,
Fortaleza - CE, C.P. 6030, 60455-760 - Brazil*

²*Instituto Federal de Educação, Ciência e Tecnologia do Ceará (IFCE),
Campus Juazeiro do Norte, 63040-000 Juazeiro do Norte - CE - Brazil*

(Dated: April 9, 2013)

Abstract

Considering quantum gravity within the framework of effective field theory, we investigated the consequences of spontaneous Lorentz violation for the gravitational potential. In particular, we focus our attention on the bumblebee models, in which the graviton couples to a vector B_μ that assumes a nonzero vacuum expectation value. The leading order corrections for the nonrelativistic potential are obtained from calculation of the scattering matrix of two scalar particles interacting gravitationally. These corrections imply a modification on the magnitude of the interaction and also add a Darwin-like term for Newton's potential.

PACS numbers: 11.30.Cp, 04.25.Nx, 12.60.-i

*Electronic address: r.v.maluf@fisica.ufc.br

†Electronic address: victor.santos@fisica.ufc.br

‡Electronic address: wilamicruz@gmail.com

§Electronic address: carlos@fisica.ufc.br

I. INTRODUCTION

A longstanding problem in theoretical physics is the conciliation between the Standard Model (SM) describing the behaviour of elementary particles and General Relativity (GR), which accounts the large scale physics dominated by gravity. With such a conciliation, both theories, which are extremely well tested, should appear as low-energy descriptions of a single and fundamental (and yet unknown) theory of quantum gravity. This framework opens the possibility for the discovery of new phenomena, not described by any of these effective theories. Unfortunately, since quantum gravity effects are relevant at energy scales of the order of the Planck mass $m_{\text{P}} \sim 1.22 \times 10^{19}$ GeV, no experimental evidence for signature of a more fundamental physics was obtained up to the current date.

Despite of the fact that Planck scale dynamics remains impossible to be accessed experimentally, a great deal of works have been performed by exploring the point of view that quantum gravity phenomena can be observed by amplification of its effects at attainable energies. One of the most interesting possibilities is the violation of Lorentz symmetry [1]. In fact, the existence of different mechanisms that bring out Lorentz-violating (LV) effects is supported in several theoretical contexts, such as loop quantum gravity [2], string theory [3], noncommutative field theories [4], and more recently in warped brane worlds [5, 6] and Horava-Lifshitz gravity [7].

The first framework to account LV in the SM was proposed by Colladay and Kostelecký [8], based on the idea of spontaneous Lorentz symmetry breaking in the context of string theory [9], known as the Standard Model Extension (SME). The SME provides a set of gauge-invariant LV tensor operators, compatible with the coordinate invariance [10], and suitable to address the CPT and Lorentz violation in physical systems. A number of interesting investigations have been developed in the different sectors of the SME. The CPT-even gauge sector was first examined by Kostelecký and Mewes [11], with the attainment of upper bound of 1 part in 10^{37} (using birefringence data). This sector was also addressed in connection with its classical solutions [12], consistency aspects [13] and fermion/photon interactions [14, 15]. More recently, new works have proposed LV scenarios endowed with higher dimensional operators, with new interesting results [16, 17]. Higher dimensional operators can be considered in terms of nonminimal interactions as well. A CPT-odd nonminimal coupling for fermions was first regarded in Ref. [18], with some recent developments [19]. Very recently,

an analogue CPT-even nonminimal coupling for fermions, embracing the K_F gauge tensor of the SME, was proposed and discussed both in relativistic and nonrelativistic scenarios [20].

Another relevant SME sector much addressed in the latest years is the gravitational one. The SME accommodates both explicit symmetry breaking as well as spontaneous breaking. However, when one focus in its gravitational sector, one notices that the explicit violation is incompatible with geometrical identities like the Bianchi identity, reason by which one should work with spontaneous breakings to address LV within the gravitational sector [21]. A general treatment of spontaneous local Lorentz and diffeomorphism violation for the gravitational sector of the SME was first addressed in Refs. [22, 23]. In these papers, it is supposed that tensor fields acquire nonzero vacuum expectation values (VEV), breaking these symmetries spontaneously. It was then shown that the corresponding linearized effective equations can be used to study the post-Newtonian effects in a series of gravitational systems [24–26]. It is worth mentioning that a discussion for alternative ways to introduce Lorentz violation in gravity was considered in [27].

In this work we investigate low-energy effects of Lorentz violation in the context of the gravitational sector of SME. More precisely, we use a particular choice in which the spontaneous Lorentz violation comes from the dynamics of a single vector field B_μ , the so called bumblebee model [21, 23], to calculate quantum corrections for the gravitational potential. In the weak-field approximation of this model, we show that the introduction of an uncharged scalar field, coupled with the gravitational field, leads to corrections to the classical Newtonian potential, including a Darwin term. The theoretical modifications here obtained are used to set up stringent bounds on the magnitude of the breaking parameter. Along all this work we shall use the spacetime signature $(+ - - -)$ and adopt the following definition for the Ricci tensor: $R_{\mu\nu} = \partial_\sigma \Gamma_{\mu\nu}^\sigma - \partial_\nu \Gamma_{\mu\sigma}^\sigma + \Gamma_{\sigma\lambda}^\lambda \Gamma_{\mu\nu}^\sigma - \Gamma_{\sigma\nu}^\lambda \Gamma_{\mu\lambda}^\sigma$, where $\Gamma_{\mu\nu}^\lambda = \frac{1}{2}g^{\lambda\sigma}(\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu})$. All quantities are expressed in natural units ($\hbar = c = \epsilon_0 = 1$), in which the gravitational constant is $G_N = 6.707 \times 10^{-57} \text{eV}^{-2}$. Moreover, tensors are symmetrized with unit weight, i.e., $A_{(\mu\nu)} = \frac{1}{2}(A_{\mu\nu} + A_{\nu\mu})$.

The structure of the paper is as follows. Sec. II is devoted to discuss the theoretical model, introducing the general action including a LV term, and then restricting to spontaneous LV. In Sec. III, we perform the weak-field approximation and calculate the LV-corrected propagator. In Sec. IV, we introduce the coupling with a matter field and obtain the

nonrelativistic potential for two bosons interacting gravitationally, via a scattering process. In Sec. V we present some constraints and limits to our results. Finally, we present our final remarks in Sec. VI.

II. THE THEORETICAL MODEL

The simplest gravity model involving Lorentz-violating terms that combine tensor fields, responsible for the spontaneous local Lorentz breaking, with the gravitational field in (3+1)-dimensional Riemann spacetime, is given by the action

$$S = S_{\text{EH}} + S_{\text{LV}} + S_{\text{matter}}. \quad (1)$$

The first piece in the above equation represents the usual Einstein-Hilbert action, defined by

$$S_{\text{EH}} = \int d^4x \sqrt{-g} \frac{2}{\kappa^2} (R - 2\Lambda), \quad (2)$$

where g denotes the determinant of the metric field $g_{\mu\nu}$, R is the Ricci scalar, Λ is the cosmological constant and $\kappa^2 = 32\pi G_N$ is the gravitational coupling. Since our main goal is to examine the effects of the Lorentz-violating on the nonrelativistic gravitational potential, we can disregard the implications of Λ , assuming it equal to zero hereafter.

The second piece in Eq. (1) represents the gravitational sector for the minimal SME and contains the coefficients for Lorentz violation, coupled to the Riemann, Ricci, and scalar curvatures, in the following form (see, f.e., [23]):

$$S_{\text{LV}} = \int d^4x \sqrt{-g} \frac{2}{\kappa^2} (uR + s^{\mu\nu} R_{\mu\nu} + t^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta}), \quad (3)$$

where u , $s^{\mu\nu}$ and $t^{\mu\nu\alpha\beta}$ are dynamical tensor fields with zero mass dimension and the same symmetries of their geometric tensors associated. This action is assumed to be invariant under general coordinate transformations and the local Lorentz violation must be achieved through a Higgs-like mechanism.

The last term on the right side of Eq. (1) takes into account the matter-gravity couplings, which in principle should include all fields of the standard model as well as possible interactions with coefficients u , $s^{\mu\nu}$ and $t^{\mu\nu\alpha\beta}$. However, we will focus our attention on the possible effects produced by the action (3), restricting ourselves to the case where the ordinary matter only interacts with the gravitational field. Further details about these effects

in the context of Lorentz-violation involving the matter sector of the SME can be seen in Ref. [28].

Next, let us consider the particular case when $t^{\mu\nu\alpha\beta} = 0$. The coefficients u and $s^{\mu\nu}$ have 10 degrees of freedom (the trace of $s^{\mu\nu}$ could be absorbed in the scalar coefficient u) that may be described by an effective field theory involving a single vector field B_μ , whose dynamics is determined by the following action

$$S_B = \int d^4x \sqrt{-g} \left[-\frac{1}{4} B_{\mu\nu} B^{\mu\nu} + \sigma B^\mu B^\nu R_{\mu\nu} - V(B_\mu B^\mu \mp b^2) \right], \quad (4)$$

where $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$, σ is a dimensionless coupling constant and b^2 is a positive constant that sets the VEV for B_μ . The potential $V(x)$ triggers the spontaneous breakdown of both Lorentz and diffeomorphism symmetries, such that its minimum occurs at $g^{\mu\nu} B_\mu B_\nu \pm b^2 = 0$, i.e., when B_μ and $g_{\mu\nu}$ acquire nonzero vacuum expectation values. This theory is a particular case of the so-called bumblebee models and were initially evaluated in the context of string theory [9]. Furthermore, we note that for $\sigma = 0$ the action for the bumblebee field becomes $U(1)$ gauge invariant and the potential V also spontaneously breaks this symmetry.

The correspondence between the action (3) and the bumblebee model (4) is obtained through the relations [23]:

$$\begin{aligned} u &= \frac{1}{4} \xi B^\alpha B_\alpha, \\ s^{\mu\nu} &= \xi B^\mu B^\nu - \frac{1}{4} \xi g^{\mu\nu} B^\alpha B_\alpha, \\ t^{\mu\nu\alpha\beta} &= 0, \end{aligned} \quad (5)$$

where for convenience we write $\sigma = (2\xi/\kappa^2)$, so that the mass dimension of the bumblebee field and the coupling constant are, respectively: $[B^\mu] = 1$, $[\xi] = -2$.

III. WEAK-FIELD APPROXIMATION AND THE GRAVITON PROPAGATOR

To investigate the effects of gravity-bumblebee coupling on the graviton dynamics, we split the dynamical fields into the vacuum expectation values and the quantum fluctuations:

$$\begin{aligned} g_{\mu\nu} &= \eta_{\mu\nu} + \kappa h_{\mu\nu}, \\ B_\mu &= b_\mu + \tilde{B}_\mu, \\ B^\mu &= b^\mu + \tilde{B}^\mu - \kappa b_\nu h^{\mu\nu}, \end{aligned} \quad (6)$$

where $h_{\mu\nu}$ and $\tilde{B}_\mu$ represent small perturbations around the Minkowski background and a constant vacuum value b_μ , respectively. The vector b_μ is the local Lorentz violation coefficient associated to the bumblebee field.

Varying the action (4) with respect to B_μ , we obtain the equation of motion for the bumblebee field:

$$\frac{1}{\sqrt{-g}}\partial^\mu \{\sqrt{-g}B_{\mu\nu}\} - 2V'B_\nu + 2\sigma B^\mu R_{\mu\nu} = 0, \quad (7)$$

where the prime on V means differentiation with respect to the argument.

Following the ideas described in Ref [23], we may employ the expansions defined in Eq. (6), and assume for $V(x)$ the smooth quadratic form

$$V = \frac{\lambda}{2} (B^\mu B_\mu \mp b^2)^2, \quad (8)$$

so that the linearized version of the equation of motion (7) can be written as

$$(\square\eta_{\mu\nu} - \partial_\mu\partial_\nu - 4\lambda b_\mu b_\nu)\tilde{B}^\mu = -2\lambda\kappa b_\nu b_\alpha b_\beta h^{\alpha\beta} - 2\sigma b^\alpha R_{\alpha\nu}, \quad (9)$$

with $\square \equiv \partial^2$. In this expression, $R_{\mu\nu}$ shall be understood as being in its linearized form. Also, for simplicity b_μ is adopted as a timelike vector, such that $b^\mu b_\mu = +b^2$. Applying the Green's function method, the solution to Eq. (9) is straightforward, leading in momentum space to the the following expression

$$\tilde{B}^\mu = \frac{\kappa p^\mu b_\alpha b_\beta h^{\alpha\beta}}{2b \cdot p} + \frac{2\sigma b_\alpha R^{\alpha\mu}}{p^2} - \frac{2\sigma p^\mu b_\alpha b_\beta R^{\alpha\beta}}{p^2 b \cdot p} + \frac{\sigma p^\mu R}{4\lambda b \cdot p} - \frac{\sigma b^\mu R}{p^2} + \frac{\sigma p^\mu b^2 R}{p^2 b \cdot p}, \quad (10)$$

with $b \cdot p = b_\mu p^\mu$, $p^2 = p \cdot p = p_\mu p^\mu$.

By substituting this solution into the action (3), with the help of the relations defined by Eqs. (5) and (6) in a suitable order, we are able to determine the modifications yielded by the nonzero vacuum expectation value b_μ on the kinetic terms of the graviton field. Therefore, it is necessary to expand the bumblebee-graviton interaction $\mathcal{L}_{LV}$ up to second order in $h_{\mu\nu}$, as it follows below:

$$\begin{aligned} \mathcal{L}_{LV} &= \sigma\sqrt{-g}B^\mu B^\nu R_{\mu\nu} \\ &= \sigma \left[b_\mu b_\nu R^{\mu\nu}(h^2) + 2b_\mu \tilde{B}_\nu R^{\mu\nu}(h) + \frac{1}{2}\kappa h^\alpha{}_\alpha b_\mu b_\nu R^{\mu\nu}(h) \right] + \mathcal{O}(h^3), \end{aligned} \quad (11)$$

where the order in $h_{\mu\nu}$ at the Ricci tensors is explicitly indicated. Replacing $\tilde{B}^\mu$ and grouping

the terms conveniently, we obtain

$$\begin{aligned}
\mathcal{L}_{\text{LV}} = & \xi \left[p^2 b_\mu b_\nu h^{\mu\nu} h^\alpha_\alpha + \frac{1}{2} (b \cdot p)^2 (h^\alpha_\alpha)^2 \right. \\
& - \frac{1}{2} (b \cdot p)^2 h^{\mu\nu} h_{\mu\nu} + p^2 b_\mu b_\nu h^{\mu\alpha} h^\nu_\alpha - (b_\mu b_\nu p_\alpha p_\beta + b_{(\mu} p_{\nu)}) h^{\mu\nu} h^{\alpha\beta} \left. \right] \\
& + \frac{4\xi^2}{\kappa^2} \left[\left(-2p^2 b_\mu b_\nu - 2b^2 p_\mu p_\nu + 4b \cdot p b_{(\mu} p_{\nu)} - \frac{p^2 p_\mu p_\nu}{4\lambda} \right) h^{\mu\nu} h^\alpha_\alpha \right. \\
& + \left(2b_\mu b_\nu p_\alpha p_\beta - b_{(\mu} p_{\nu)} b_{(\alpha} p_{\beta)} + \frac{b^2 p_\mu p_\nu p_\alpha p_\beta}{p^2} - \frac{2b \cdot p p_\mu p_\nu b_{(\alpha} p_{\beta)}}{p^2} + \frac{p_\mu p_\nu p_\alpha p_\beta}{4\lambda} \right) h^{\mu\nu} h^{\alpha\beta} \\
& \left. + \left(b^2 p^2 - (b \cdot p)^2 + \frac{p^4}{4\lambda} \right) (h^\alpha_\alpha)^2 + \left(p^2 b_\mu b_\nu - 2b \cdot p b_{(\mu} p_{\nu)} + \frac{(b \cdot p)^2 p_\mu p_\nu}{p^2} \right) h^{\mu\lambda} h^\nu_\lambda \right] + \mathcal{O}(h^3),
\end{aligned} \tag{12}$$

with $\sigma = (2\xi/\kappa^2)$, as previously defined. The above expression can be rewritten to position space and combined with the expanded Einstein-Hilbert Lagrangian

$$\mathcal{L}_{\text{EH}} = \partial h^{\mu\nu} \partial_\alpha h^\alpha_\nu - \partial_\mu h^{\mu\nu} \partial_\nu h + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{1}{2} \partial_\alpha h^{\mu\nu} \partial^\alpha h_{\mu\nu} + \mathcal{O}(h^3), \tag{13}$$

with $h \equiv h^\lambda_\lambda$. We add one convenient gauge fixing term,

$$\mathcal{L}_{\text{gf}} = -(\partial_\mu h^{\mu\nu} - \frac{1}{2} \partial^\nu h)^2, \tag{14}$$

to yield the effective Lagrangian, which we need to consider in order to obtain the modified graviton propagator. Then, in the $\mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{gf}} + \mathcal{L}_{\text{LV}}$, the kinetic term for the graviton field becomes

$$\mathcal{L}_{\text{kin}} = -\frac{1}{2} h^{\mu\nu} \hat{\mathcal{O}}_{\mu\nu, \alpha\beta} h^{\alpha\beta}, \tag{15}$$

where the operator $\hat{\mathcal{O}}_{\mu\nu, \alpha\beta}$ is separated in two pieces

$$\hat{\mathcal{O}}_{\mu\nu, \alpha\beta} = \hat{\mathcal{K}}_{\mu\nu, \alpha\beta} + \hat{\mathcal{V}}_{\mu\nu, \alpha\beta}, \tag{16}$$

such that $\hat{\mathcal{K}}_{\mu\nu, \alpha\beta}$ is the usual quadratic form,

$$\hat{\mathcal{K}}_{\mu\nu, \alpha\beta} = \frac{1}{2} (\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha} - \eta_{\mu\nu} \eta_{\alpha\beta}) (-\partial^2), \tag{17}$$

while $\hat{\mathcal{V}}_{\mu\nu, \alpha\beta}$ encloses the terms that contain the Lorentz-violating Lagrangian $\mathcal{L}_{\text{LV}}$.

The graviton propagator is defined by

$$\langle 0 | T [h_{\mu\nu}(x) h_{\alpha\beta}(y)] | 0 \rangle = D_{\mu\nu, \alpha\beta}(x - y), \tag{18}$$

where $D_{\mu\nu,\alpha\beta}$ is the operator that satisfies the Green's equation, given as

$$\hat{O}^{\mu\nu}_{\lambda\sigma} D^{\lambda\sigma,\alpha\beta}(x-y) = i\mathcal{I}^{\mu\nu,\alpha\beta} \delta^4(x-y), \quad (19)$$

with $\mathcal{I}^{\mu\nu,\alpha\beta} = \frac{1}{2} (\eta^{\mu\alpha} \eta^{\nu\beta} + \eta^{\mu\beta} \eta^{\nu\alpha})$. Thus, the exact graviton propagator is evaluated by inverting (16), finding a closed operator algebra composed by a set of appropriated projectors. However, in order to avoid problems concerning the non-physical modes induced by the higher derivative terms, we take advantage of the fact that b_μ is small. We therefore use the conventional graviton propagator in the gauge given by Eq. (14) and treat the Lorentz-violating term in Eq. (16) as a perturbative insertion [29]. This is accomplished by means of the following matricial identity:

$$\frac{1}{A+B} = \frac{1}{A} - \frac{1}{A} B \frac{1}{A+B} = \frac{1}{A} - \frac{1}{A} B \frac{1}{A} + \frac{1}{A} B \frac{1}{A} B \frac{1}{A+B} = \dots \quad (20)$$

The operator $\hat{\mathcal{K}}$ can easily be inverted and the conventional graviton propagator is then written in the momentum space as

$$D_0^{\mu\nu,\alpha\beta}(q) = \frac{i}{2} \frac{\eta^{\mu\alpha} \eta^{\nu\beta} + \eta^{\mu\beta} \eta^{\nu\alpha} - \eta^{\mu\nu} \eta^{\alpha\beta}}{q^2 + i\epsilon}. \quad (21)$$

After lengthy contraction operations of indices, we are ready to give the explicit form of $D^{\mu\nu,\alpha\beta} = D_0^{\mu\nu,\alpha\beta} + D_{\text{LV}}^{\mu\nu,\alpha\beta}$ up to second order in b_μ , which reads as below:

$$\begin{aligned} \left(D_{\text{LV}}^{\mu\nu,\alpha\beta} \right)_\xi = & i\xi \left[b^2 \left(\frac{g^{\alpha\beta} g^{\mu\nu}}{q^2} + \frac{q^\alpha q^\beta g^{\mu\nu}}{q^4} \right) \right. \\ & + \frac{(b \cdot q)^2 (g^{\alpha\beta} g^{\mu\nu} - g^{\alpha\nu} g^{\beta\mu} - g^{\alpha\mu} g^{\beta\nu})}{2q^4} \\ & + \frac{b \cdot q (b^\beta q^\alpha g^{\mu\nu} + b^\alpha q^\beta g^{\mu\nu} + b^\nu q^\mu g^{\alpha\beta} + b^\mu q^\nu g^{\alpha\beta})}{2q^4} \\ & + \left(\frac{b^\alpha b^\mu g^{\beta\nu} + b^\beta b^\mu g^{\alpha\nu} + b^\alpha b^\nu g^{\beta\mu} + b^\beta b^\nu g^{\alpha\mu} - 2b^\alpha b^\beta g^{\mu\nu} - 4b^\mu b^\nu g^{\alpha\beta}}{2q^2} \right. \\ & \left. \left. - \frac{4b^\mu b^\nu q^\alpha q^\beta + b^\beta b^\nu q^\alpha q^\mu + b^\alpha b^\nu q^\beta q^\mu + b^\beta b^\mu q^\alpha q^\nu + b^\alpha b^\mu q^\beta q^\nu}{2q^4} \right) \right], \quad (22) \end{aligned}$$

$$\begin{aligned}
\left(D_{\text{LV}}^{\mu\nu,\alpha\beta}\right)_{\xi^2} = & \frac{i\xi^2}{\kappa^2} \left[b^2 \left(\frac{12q^\mu q^\nu g^{\alpha\beta} - 12q^\alpha q^\beta g^{\mu\nu}}{q^4} + \frac{8q^\alpha q^\beta q^\mu q^\nu}{q^6} \right) + \frac{g^{\alpha\beta} g^{\mu\nu}}{2\lambda} \right. \\
& + \frac{2(b \cdot q)^2 (q^\alpha q^\mu g^{\beta\nu} + q^\beta q^\mu g^{\alpha\nu} + q^\alpha q^\nu g^{\beta\mu} + q^\beta q^\nu g^{\alpha\mu} + 2q^\mu q^\nu g^{\alpha\beta} - 2q^\alpha q^\beta g^{\mu\nu})}{q^6} \\
& + b \cdot q \left\{ \frac{10(b^\beta q^\alpha g^{\mu\nu} + b^\alpha q^\beta g^{\mu\nu} - b^\nu q^\mu g^{\alpha\beta} - b^\mu q^\nu g^{\alpha\beta})}{q^4} + \frac{8(b^\beta q^\alpha q^\mu q^\nu + b^\alpha q^\beta q^\mu q^\nu)}{q^6} \right. \\
& - \left. \frac{4(b^\mu q^\alpha g^{\beta\nu} - b^\mu q^\beta g^{\alpha\nu} - b^\nu q^\alpha g^{\beta\mu} - b^\nu q^\beta g^{\alpha\mu})}{q^4} \right\} - \frac{q^\alpha q^\beta g^{\mu\nu}}{q^2 \lambda} + \frac{3q^\mu q^\nu g^{\alpha\beta}}{q^2 \lambda} \\
& + \left\{ \frac{2(b^\alpha b^\mu g^{\beta\nu} + b^\beta b^\mu g^{\alpha\nu} + b^\alpha b^\nu g^{\beta\mu} + b^\beta b^\nu g^{\alpha\mu} - 2b^\alpha b^\beta g^{\mu\nu} + 2b^\mu b^\nu g^{\alpha\beta})}{q^2} \right. \\
& + \left. \frac{2(8b^\mu b^\nu q^\alpha q^\beta - b^\beta b^\nu q^\alpha q^\mu - b^\alpha b^\nu q^\beta q^\mu - b^\beta b^\mu q^\alpha q^\nu - b^\alpha b^\mu q^\beta q^\nu) + \frac{2q^\alpha q^\beta q^\mu q^\nu}{\lambda}}{q^4} \right\} \Big] \quad (23)
\end{aligned}$$

where $\left(D_{\text{LV}}^{\mu\nu,\alpha\beta}\right)_\xi$ and $\left(D_{\text{LV}}^{\mu\nu,\alpha\beta}\right)_{\xi^2}$ are contributions to $D_{\text{LV}}^{\mu\nu,\alpha\beta}$ proportional to ξ and ξ^2 , respectively.

Some comments about these results are worthwhile. Taking into account the expression (5), the products b^2 , $(b \cdot q)^2$ and $(b \cdot q)b^\mu$ are first order terms in the Lorentz-violating coefficients u and $s^{\mu\nu}$. Thus, we note that the correction $\left(D_{\text{LV}}^{\mu\nu,\alpha\beta}\right)_\xi$ involves only terms to first order in u and $s^{\mu\nu}$, and do not depend on the particular form of the potential $V(x)$. However, in second order in ξ , there are terms that are not associated with the vector b_μ (they are proportional to λ^{-1}) and depend only on the coupling of this potential. In addition, the corrections to graviton propagator have poles in $q^2 = 0$, showed that in this approximation the theory is free of ghosts and tachyons. Finally, it is still important to mention that this propagator is symmetric under an indices permutation ($\mu \leftrightarrow \nu$) and ($\alpha \leftrightarrow \beta$), as it really must be. We should draw attention for other evaluations concerning the graviton propagator [30].

IV. MODIFIED NEWTON'S LAW OF GRAVITATION

In this section, we study the effects of the spontaneous Lorentz violation when we consider the tree-level modified propagator as determined previously. One of the simplest examples that we can choose to evaluate such effects, consists in the gravitational interaction of two distinguishable heavy particles described in the nonrelativistic limit by the Newtonian po-




Figure 1: The tree-level diagram of two scalar particles interacting via the exchange of a graviton.

tential. Thus, our main goal here is to determine the scattering amplitude of two massive bosons particles of spin-zero by one-graviton exchange. Once calculated the matrix amplitude in leading-order, we can take the nonrelativistic limit and compare it with the Born approximation to determine the potential modified by the nonzero vacuum expectation value b_μ .

Consider the following action for a real scalar field in curved spacetime,

$$S_{\text{matter}} = \int d^4x \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} m^2 \phi^2 \right], \quad (24)$$

which can be expanded in the weak field approximation up to first order in h . We are then left with the following Lagrangian:

$$\begin{aligned} \mathcal{L}_{\text{matter}} \approx & \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 \\ & - \frac{1}{2} \kappa h^{\mu\nu} \left[\partial_\mu \phi \partial_\nu \phi - \frac{1}{2} \eta_{\mu\nu} (\partial_\alpha \phi \partial^\alpha \phi - m^2 \phi^2) \right]. \end{aligned} \quad (25)$$

Now, let us consider the scattering process involving two scalar particles of mass m_1 and m_2 . The only Feynman diagram that contributes to this process, in lowest order, is drawn in Fig. 1, and its analytical expression can be written as

$$i\mathcal{M} = (-i\kappa)^2 V^{\mu\nu}(p_1, -k_1, m_1) D_{\mu\nu, \alpha\beta}(q) V^{\alpha\beta}(p_2, -k_2, m_2), \quad (26)$$

where $q = p_2 - k_2 = -(p_1 - k_1)$ is the momentum transfer and the vertex $V^{\mu\nu}(p, k, m)$ corresponds to the expression

$$V^{\mu\nu}(p, k, m) = -\frac{1}{2} [p^\mu k^\nu + p^\nu k^\mu - \eta^{\mu\nu} (p \cdot k + m^2)]. \quad (27)$$

Substituting the expressions defined in (18) and (27) into the scattering amplitude (26), we arrive at the sum of the two pieces:

$$i\mathcal{M} = i\mathcal{M}_0 + i\mathcal{M}_{\text{LV}}, \quad (28)$$

such that the first term is just the conventional amplitude given by [31]

$$i\mathcal{M}_0 = -\frac{i\kappa^2}{8q^2} \left[4 \{ k_1 \cdot p_1 (m_2^2 - k_2 \cdot p_2) + k_1 \cdot p_2 k_2 \cdot p_1 + k_1 \cdot k_2 p_1 \cdot p_2 \} \right. \\ \left. - 2m_1^2 \{ 4 (m_2^2 - k_2 \cdot p_2) + 2k_2 \cdot p_2 \} \right], \quad (29)$$

which is modified by $i\mathcal{M}_{\text{LV}}$, consisting of a large expression involving the possible contractions of b_μ with the external momenta.

To access the nonrelativistic limit, we take the approximation $p_{1,2} = (m_{1,2}, 0)$, $k_{1,2} = (m_{1,2}, 0)$, and $q = (0, \vec{q})$, which leads to the following result:

$$i\mathcal{M}_{\text{NR}} = \frac{i\kappa^2 m_1^2 m_2^2}{2\vec{q}^2} - \frac{i\xi \vec{b}^2 \kappa^2 m_1^2 m_2^2}{\vec{q}^2} + \frac{8i\xi^2 b_0^2 m_1^2 m_2^2}{\vec{q}^2} + \frac{i\xi^2 m_1^2 m_2^2}{2\lambda}, \quad (30)$$

where the last three terms involve the correction matrix arising from the spontaneous Lorentz breaking, so that $b^\mu = (b^0, \vec{b})$ is the constant background.

To make the connection to the Newtonian gravitational potential, we follow Ref. [32], and define the potential Fourier transformed in the nonrelativistic limit by

$$\langle f | iT | i \rangle \equiv (2\pi)^4 \delta^4(p - k) i\mathcal{M}(p_1, p_2 \rightarrow k_1, k_2) \\ \approx -(2\pi) \delta(E_p - E_k) i\tilde{V}(\vec{q}), \quad (31)$$

so that the potential in coordinate space corresponds to

$$V(\vec{x}) = \frac{1}{2m_1} \frac{1}{2m_2} \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q} \cdot \vec{x}} \tilde{V}(\vec{q}). \quad (32)$$

Finally, after inverting the equation (30), we obtain the desired result

$$V(\vec{x}) = -\frac{G_N m_1 m_2}{|\vec{x}|} + \frac{2\xi \vec{b}^2 G_N m_1 m_2}{|\vec{x}|} - \frac{\xi^2 b_0^2 m_1 m_2}{2\pi |\vec{x}|} - \frac{\xi^2 m_1 m_2}{8\lambda} \delta^{(3)}(\vec{x}). \quad (33)$$

As one can see, the first term above is the standard Newton's potential. The next two terms only scale the magnitude of interaction. However, the last term provides a nontrivial contribution, involving a Dirac delta function. This correction looks like a Darwin's term, which also appears in the electron-photon scattering in QED and it is interpreted as quantum fluctuations from the motion of the electron due to its position uncertainty. Moreover, a similar result was obtained in the context of the Horava-Lifshitz gravity in Ref. [33].

Next, we will show that the result in Eq. (33) is suitable to give constraints on the magnitude scale of the coefficients for Lorentz violation.

V. EXPERIMENTAL CONSTRAINTS: SOLAR GRAVITATIONAL DEFLECTION

As mentioned earlier, one of the effects of the non-vanishing VEV b^μ is a rescaling of the gravitational constant G_N by $\tilde{G} = G_N(1 - 2\xi\vec{b}^2 + \frac{\xi^2 b_0^2}{2\pi G_N})$. However, experiments sensitive to Lorentz violation in the matter and gauge sectors of the SME indicate that the coefficients for Lorentz violation are very small [34]. This feature is also expected in the gravitational sector of the SME, where we can use the standard experimental tests of GR to achieve stringent bounds on Lorentz violation [35]. In this context, the deflection of light by the Sun is a convenient experiment to apply our result.

The deflection angle by which an light ray is deflected by the Sun is given by (see Ref. [36] for more details.)

$$\Delta \simeq \frac{4G_N M_\odot}{d}, \quad (34)$$

where $M_\odot = 1.99 \times 10^{30}$ kg $\approx 1.11 \times 10^{66}$ eV is the mass of the Sun and d is the impact parameter which can be effectively considered as the Sun's radius $R_\odot = 6.96 \times 10^8$ m $\approx 3.53 \times 10^{15}$ (eV) $^{-1}$ (Note that $c = 1$ in our units). Using GR, for a light beam passing near the Sun, $\Delta \approx 1.75$ arcsec $\approx 8.49 \times 10^{-6}$ rad. Considering that the experimental results based on interferometric observations of the deflection of radio waves by the Sun are able to detect effects of 1 part in 10^3 approximately [37], the corrections to Eq. (34) resulting from (33) may not be larger than 10^{-9} rad.

For our initial evaluation, we consider the correction of first order in ξ , a quantity which is associated with the spatial component of b^μ . This correction is $\Delta_\xi = -8\xi\vec{b}^2 G_N M_\odot / R_\odot$, whose numerical value is $\simeq 1.68 \times 10^{-5} |\xi\vec{b}^2|$. Taking into account that the deflection angle will be undetectable if it is less than 10^{-9} , we obtain the following bound for the parameter product: $|\xi\vec{b}^2| \leq 5.95 \times 10^{-5}$.

In the case of the term that involves the timelike component of b^μ , we arrived at the following result for the gravitational deflection correction: $\Delta_{\xi^2} = 2(\xi b_0)^2 M_\odot / (\pi R_\odot)$, whose numerical value is $\simeq 2.00 \times 10^{50} (\xi b_0)^2 (\text{eV})^2$, leading to the upper bound $|\xi b_0| \leq 2.23 \times 10^{-30} (\text{eV})^{-1}$. The order of magnitude of this bound is much more restrictive than the one obtained previously, involving the component $\vec{b}$. However, this difference is expected, since the product of the parameters $\xi\vec{b}^2$ is a dimensionless quantity, its bound depends only on the accuracy in the experimental measurements considered.

VI. CONCLUSIONS

In this work, we presented the modifications produced by the spontaneous breaking of Lorentz symmetry over the Newtonian gravitational potential by means of the direct calculation of the scattering amplitude between two massive scalar particles interacting gravitationally.

Firstly, we have introduced an action to the simplest gravity model involving tensor fields, responsible for the spontaneous local Lorentz breaking, coupled with the gravitational field. To construct the gravitational sector for the minimal SME we take a particular case of the so-called bumblebee model. After that, we separate the dynamical fields into the vacuum expectation values and the quantum fluctuations to analyze the effects of gravity-bumblebee coupling on the graviton dynamics. Inserting the solution of the equation of motion for the bumblebee field in the LV action, we have determined the modifications the kinetic terms of the graviton field. Dealing these modifications in the form of a perturbative insertion, we have obtained a corrected propagator for which ghosts and tachyons are not present.

As a result, we observed at first order in LV coupling ξ , a modification in the coefficients which can be absorbed in the Newton's constant G_N and hence provide experimental bounds to the LV parameter, based on the accurate measures of the deflection of light by the Sun. In second order in ξ , we verify the appearance of a Darwin-like correction term, independent of the VEV b_μ of the bumblebee field, reflecting the effect of the spontaneous symmetry breaking potential $V(B_\mu B^\mu \mp b^2)$. This result corroborates the fact that at short distances, where the gravitational field is intense, the local Lorentz symmetry might be violated.

Acknowledgments

This work was partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Pesquisas (CNPq) and Fundação Cearense de apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP).

[1] D. Mattingly, Living Rev. Rel. **8**, 5 (2005).

- [2] J. Alfaro, H. A. Morales-Tecotl and L. F. Urrutia, Phys. Rev. Lett. **84**, 2318 (2000); J. Alfaro, H. A. Morales-Tecotl and L. F. Urrutia, Phys. Rev. D **65**, 103509 (2002).
- [3] N. E. Mavromatos, PoS QG-PH, **027** (2007).
- [4] S. M. Carroll, J. A. Harvey, V. A. Kostelecký, C. D. Lane, and T. Okamoto, Phys. Rev. Lett. **87**, 141601 (2001).
- [5] T. G. Rizzo, JHEP **0509** 036 (2005); T. G. Rizzo, JHEP **1011** 156 (2010).
- [6] V. Santos and C. A. S. Almeida, Phys. Lett. B **718**, 1114 (2013).
- [7] M. Pospelov and Y. Shang, Phys. Rev. D **85**, 105001 (2012).
- [8] D. Colladay and V. A. Kostelecký, Phys. Rev. D **55**, 6760 (1997); Phys. Rev. D **58**, 116002 (1998).
- [9] V. A. Kostelecký and S. Samuel, Phys. Rev. D **39**, 683 (1989).
- [10] V. A. Kostelecký and R. Lehnert, Phys. Rev. D **63**, 065008 (2001).
- [11] V. A. Kostelecký and M. Mewes, Phys. Rev. Lett. **87**, 251304 (2001); Phys. Rev. D **66**, 056005 (2002); Phys. Rev. Lett. **97**, 140401 (2006).
- [12] Q. G. Bailey and V. A. Kostelecký, Phys. Rev. D **70**, 076006 (2004); R. Casana, M.M. Ferreira Jr, C. E. H. Santos, Phys. Rev. D **78**, 105014 (2008); R. Casana, M.M. Ferreira Jr, A. R. Gomes, P. R. D. Pinheiro, Eur. Phys. J. C **62**, 573 (2009).
- [13] R. Casana, M.M. Ferreira Jr, A. R. Gomes, P. R. D. Pinheiro, Phys. Rev. D **80**, 125040 (2009); R. Casana, M.M. Ferreira Jr, A. R. Gomes, F. E. P. dos Santos, Phys. Rev. D **82**, 125006 (2010); F.R. Klinkhamer, M. Schreck, Nucl. Phys. B **848**, 90 (2011); M. Schreck, Phys. Rev. D **86**, 065038 (2012).
- [14] F. R. Klinkhamer and M. Risse, Phys. Rev. D **77**, 016002 (2008); **77**, 117901 (2008); F. R. Klinkhamer and M. Schreck, Phys. Rev. D **78**, 085026 (2008); B. Charneski, M. Gomes, T. Mariz, J. R. Nascimento and A. J. da Silva, Phys. Rev. D **79**, 065007 (2009).
- [15] B. Altschul, Nucl. Phys. B **796**, 262 (2008); B. Altschul, Phys. Rev. Lett. **98**, 041603 (2007); C. Kaufhold and F.R. Klinkhamer, Phys. Rev. D **76**, 025024 (2007).
- [16] V. A. Kostelecký and M. Mewes, Phys. Rev. D **80**, 015020 (2009); M. Cambiaso, R. Lehnert, and R. Potting, Phys. Rev. D **85**, 085023 (2012); M. Mewes, Phys. Rev. D **85**, 116012 (2012). B. Agostini, F. A. Barone, F. E. Barone, P. Gaete, J. A. Helayël-Neto, Phys. Lett. B **708**, 212 (2012).
- [17] R. C. Myers and M. Pospelov, Phys. Rev. Lett. **90**, 211601 (2003); P.A. Bolokhov and M.

- Pospelov, Phys. Rev. D **77**, 025022 (2008); C.M. Reyes, L. F. Urrutia, and J. D. Vergara, Phys. Rev. D **78**, 125011 (2008); C.M. Reyes, Phys. Rev. D **80**, 105008 (2009); **82**, 125036 (2010); J. Lopez-Sarrion and C. M. Reyes, Eur. Phys. J. C **72**, 2150 (2012); C.M. Reyes, L. F. Urrutia, and J. D. Vergara, Phys. Lett. B **675**, 336 (2009).
- [18] H. Belich, T. Costa-Soares, M. M. Ferreira Jr., J. A. Helayël-Neto, Eur. Phys. J. C **41**, 421 (2005); H. Belich, T. Costa-Soares, M. M. Ferreira, Jr., J. A. Helayël-Neto, and M. T. D. Orlando, Phys. Lett. B **639**, 675 (2006).
- [19] H. Belich, E. O. Silva, M. M. Ferreira Jr. and M. T. D. Orlando, Phys. Rev. D **83**, 125025 (2011); B. Charneski, M. Gomes, R. V. Maluf and A. da Silva, Phys. Rev. D **86**, 045003 (2012); G. Gazzola, H. G. Fagnoli, A. P. Baeta Scarpelli, M. Sampaio, and M. C. Nemes, J. Phys. G **39**, 035002 (2012); A. P. Baeta Scarpelli, J. Phys. G **39**, 125001 (2012); E. O. Silva and F. M. Andrade, EPL **101**, 51005 (2013).
- [20] R. Casana, M. M. Ferreira, R. V. Maluf and F. E. P. dos Santos, Phys. Rev. D **86**, 125033 (2012); R. Casana, M. M. Ferreira, Jr., E. Passos, F. E. P. dos Santos and E. O. Silva, Phys. Rev. D **87**, 047701 (2013); R. Casana, M. M. Ferreira Jr, R. V. Maluf, F. E. P. dos Santos, arXiv:1302.2375 [hep-th].
- [21] J. W. Moffat, Int. J. Mod. Phys. D **12**, 1279 (2003); V. A. Kostelecký, Phys. Rev. D **69**, 105009 (2004).
- [22] R. Bluhm and V. A. Kostelecký, Phys. Rev. D **71**, 065008 (2005).
- [23] Q. G. Bailey and V. A. Kostelecký, Phys. Rev. D **74**, 045001 (2006).
- [24] R. Bluhm, Shu-Hong Fung and V. A. Kostelecký, Phys. Rev. D **77**, 065020 (2008).
- [25] V. A. Kostelecký and R. Potting, Phys. Rev. D **79**, 065018 (2009).
- [26] B. Altschul, Q. G. Bailey and V. A. Kostelecký, Phys. Rev. D **81**, 065028 (2010).
- [27] A. F. Ferrari, M. Gomes, J. R. Nascimento, E. Passos, A. Yu. Petrov, A. J. da Silva, Phys. Lett. B **652**, 174 (2007).
- [28] V. A. Kostelecký and J. D. Tasson, Phys. Rev. D **83**, 016013 (2011).
- [29] W. F. Chen and G. Kunstatter, Phys. Rev. D **62**, 105029 (2000); C. D. Carone, M. Sher and M. Vanderhaeghen, Phys. Rev. D **74**, 077901 (2006); A. Ferrero and B. Altschul, Phys. Rev. D **84**, 065030 (2011).
- [30] J. L. Boldo, J. A. Helayël-Neto, L.M. de Moraes, C. A. G. Sasaki, V. J. V. Otoyá, Phys. Lett. B **689**, 112 (2010); C. A. Hernaski, A. A. Vargas-Paredes, J. A. Helayël-Neto, Phys. Rev. D

- 80**, 124012 (2009).
- [31] J. F. Donoghue, Phys. Rev. D **50**, 3874 (1994).
 - [32] N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, Phys. Rev. D **67**, 084033 (2003).
 - [33] F. S. Bemfica and M. Gomes, Phys. Rev. D **84**, 084022 (2011).
 - [34] V. A. Kostelecký, R. Lehnert, and M. Perry, Phys. Rev. D **68**, 123511 (2003); R. Bluhm, Lect. Notes Phys. **702**, 191 (2006).
 - [35] Floyd W. Stecker, Astroparticle Phys. **35**, 95 (2011); V. A. Kostelecký, N. Russell, Rev. Mod. Phys. **83**, 11-31 (2011).
 - [36] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*, publishing as Addison Wesley, Person Education, United States of America, 2004, pp. 285-293.
 - [37] D. E. Lebach, B.E. Corey, I. I. Shapiro, M. I. Ratner, J. C. Webber, A. E. E. Rogers, J. L. Davis, and T. A. Herring, Phys. Rev. Lett. **75**, 1439 (1995); B. Bertotti, L. Iess, and P. Tortora, Nature (London) **425**, 374 (2003); S. S. Shapiro, J. L. Davis, D. E. Lebach, and J. S. Gregory, Phys. Rev. Lett. **92**, 121101 (2004).